arXiv:0911.2814v3 [math.AG] 24 Nov 2009

A_∞-ALGEBRA OF AN ELLIPTIC CURVE AND EISENSTEIN SERIES

ALEXANDER POLISHCHUK

Abstract. We compute explicitly the A_∞-structure on the algebra $\mathrm{Ext}^*(\mathcal{O}_C \oplus L, \mathcal{O}_C \oplus L)$, where L is a line bundle of degree 1 on an elliptic curve C. The answer involves higher derivatives of Eisenstein series.

Introduction

The bounded derived category $D^b(X)$ of coherent sheaves is an important invariant of an algebraic variety X (see [1] for a survey). This category can be described in a purely algebraic way if one has a *generator*, i.e., an object G generating $D^b(X)$ as a triangulated category. With such an object one can associate a graded algebra $E_G = \oplus_{n\in\mathbb{Z}} \mathrm{Ext}^n(G,G)$. However, in order to recover $D^b(X)$ from E_G one has to take into account certain higher products which fit together into a structure of an A_∞-*algebra* on E_G (see [3] for an introduction into A_∞-algebras). Namely, one can realize E_G as the cohomology of a dg-algebra and then apply a general algebraic construction that gives an A_∞-structure on such cohomology (see [7]). This A_∞-structure is minimal in the sense that $m_1 = 0$, and is canonical up to A_∞-equivalence. Now the category $D^b(X)$ can be shown to be equivalent to the derived category of perfect A_∞-modules over A (see [4], Thm. 3.1 or [6], 7.6).

Thus, it is of interest to compute explicitly higher products on algebras of the form $\mathrm{Ext}^*(G,G)$ as above. In this paper we solve this problem in the case when X is a complex elliptic curve and $G = \mathcal{O}_X \oplus L$, where L is a line bundle of degree 1. Namely, we compute the A_∞-structure arising from the harmonic representatives (with respect to natural metrics) in the Dolbeault complex computing $\mathrm{Ext}^*(G,G)$. The resulting formulas involve higher derivatives of Eisenstein series (see Theorem 2.5.1). More precisely, we have to use the well-known non-holomorphic (but modular) modification e_2^* of the standard Eisenstein series e_2 along with all the higher Eisenstein series e_{2k}, $k \geq 2$.

It is interesting that Eisenstein series appear not in their usual form but rather as some rapidly decreasing series, similar to those considered in [9] (see Theorem 1.2.1). The A_∞-constraint gives rise to some quadratic relations involving derivatives of Eisenstein series, some of them well known (see Proposition 2.6.1).

Acknowledgment. I am grateful to Alexandr Usnich and Dmytro Shklyarov for helpful discussions. This paper was written during the stay at the Institut des Hautes Études Scientifiques. I'd like to thank this institution for hospitality and support.

Supported in part by NSF grant.

1. Eisenstein series

1.1. **Definitions.** Let us recall some basic definitions and facts (see [10], ch. III). We consider C^∞-functions $F(\omega_1, \omega_2)$ on the space of all oriented bases of $\mathbb{C} = \mathbb{R}^2$. Recall that such a function is said to be of weight $k \in \mathbb{Z}$ if

$$F(\lambda\omega_1, \lambda\omega_2) = \lambda^{-k} F(\omega_1, \omega_2).$$

Such F is called modular (with respect to $\mathrm{SL}(2, \mathbb{Z})$) if it is invariant with respect to $\mathrm{SL}(2, \mathbb{Z})$ base changes of (ω_1, ω_2) and $F(1, \tau) = f(e^{2\pi i \tau})$, where $f(q)$ is meromorphic at $q = 0$. The Eisenstein series e_{2k} for $k \geq 2$ is defined by

$$e_{2k}(\omega_1, \omega_2) = \sum_{\omega \in L \setminus \{0\}} \frac{1}{\omega^{2k}}$$

where $L = \mathbb{Z}\omega_1 + \mathbb{Z}\omega_2$. The function e_{2k} is modular of weight $2k$. One can also consider the analogous series for $k = 1$ using Eisenstein's summation rule:

$$e_2(\omega_1, \omega_2) = \sum_m \sum_{n; n \neq 0 \text{ if } m=0} \frac{1}{(m\omega_2 + n\omega_1)^2}.$$

The function e_2 is not modular but admits a simple non-holomorphic correction that makes it $\mathrm{SL}(2, \mathbb{Z})$-invariant. Namely, let us set

$$e_2^*(\omega_1, \omega_2) = e_2(\omega_1, \omega_2) - \frac{\pi}{a(L)} \frac{\bar{\omega_1}}{\omega_1},$$

where $a(L) = \mathrm{Im}(\overline{\omega}_1 \omega_2)$ is the area of $\mathbb{C}/L$. Then $e_2^*(\omega_1, \omega_2)$ is $\mathrm{SL}(2, \mathbb{Z})$-invariant (and of weight 2). For convenience we set also $e_{2k}^* = e_{2k}$ for $k \geq 2$. Eisenstein series appear as coefficients in the expansion of the Weierstrass zeta-function (cf. [10], ch. III, formula (9)):

$$\zeta(z; \omega_1, \omega_2) = \frac{1}{z} - \sum_{k \geq 2} e_{2k}(\omega_1, \omega_2) z^{2k-1}. \tag{1.1.1}$$

Following [2], sec. 1.5, we consider the normalized Weil operator

$$W = -\frac{\pi}{a(L)} (\overline{\omega}_1 \frac{\partial}{\partial \omega_1} + \overline{\omega_2} \frac{\partial}{\partial \omega_2}).$$

This operator is $\mathrm{SL}(2, \mathbb{Z})$-invariant and is of weight two. Slightly modifying the definition in [10], ch.VI, for a pair of integers $b > a \geq 0$ of different parity we set

$$g_{a,b} = g_{b,a} = (b-a)! W^a (e_{b-a+1}^*).$$

Note that $(a(L)/\pi)^a \cdot g_{a,b}$ differs by a rational factor from Weil's $e_{a,b+1}^*$. As shown in [10], sec. VI.5, $g_{a,b}$ is a polynomial in $e_2^*, e_4, \ldots, e_{a+b+1}$ with rational coefficients.

1.2. **Presentation by rapidly decreasing series.** For $m, n \in \mathbb{Z}$ and a lattice $L \subset \mathbb{C}$ we set

$$f_{m,n}(L) = (\frac{\pi}{a(L)})^m \cdot \sum_{\omega \in L \setminus \{0\}} \frac{\bar{\omega}^m}{\omega^n} \exp(-\frac{\pi}{a(L)} |\omega|^2).$$

Note that $f_{m,n} = 0$ unless $m + n$ is even (as one can see making the substitution $\omega \mapsto -\omega$).

Theorem 1.2.1. *For $n = 2k \geq 2$ one has*

$$e_n^* = \frac{2}{(n-1)!} f_{n-1,1} + \sum_{m=2}^{n} \frac{1}{(n-m)!} f_{n-m,m} = \sum_{\omega \neq 0} \frac{P_{n-1}(\frac{\pi}{a(L)}|\omega|^2)}{\omega^n} \exp(-\frac{\pi}{a(L)}|\omega|^2),$$

where

$$P_{n-1}(z) = \frac{2z^{n-1}}{(n-1)!} + \sum_{m=2}^{n} \frac{z^{n-m}}{(n-m)!}.$$

Proof. This follows easily from Theorem 1 of [9] stating that

$$\zeta(z; \omega_1, \omega_2) - z_1\eta_1 - z_2\eta_2 = \sum_{\omega \in L} \frac{\exp(-\frac{\pi}{a(L)}|\omega + z|^2)}{\omega + z} - \sum_{\omega \in L \setminus \{0\}} \frac{\exp(-\frac{\pi}{a(L)}|\omega|^2 + 2\pi i \frac{\mathrm{Im}(\overline{\omega} z)}{a(L)})}{\omega},$$

where $z = z_1\omega_1 + z_2\omega_2$ with $z_1, z_2 \in \mathbb{R}$. Indeed, we can use the expansion (1.1.1) to check the assertion for $n \geq 4$: one has to subtract $1/z$ from both parts of the above identity, apply $(\frac{\partial}{\partial z})^{n-1}$ and then evaluate at $z = 0$. The case $n = 2$ is slightly different: we again subtract $1/z$ from both parts, then apply $\frac{\partial}{\partial z_1}$ (taking (z_1, z_2) as independent variables) and evaluate at $z = 0$. The required formula follows the fact that $e_2 = -\eta_1/\omega_1$ (see e.g., [2], sec. 1.2). $\square$

Remark. The fact that e_n^* is holomorphic in (ω_1, ω_2) for $n > 2$ is equivalent to the identity

$$2f_{n-1,-1} = (n-1)f_{n-2,0}.$$

For $n = 2$ we have instead $2f_{1,-1} = f_{0,0} + 1$. These identities can be derived from the Poisson summation formula and Fourier self-duality of $\exp(-\frac{\pi}{a(L)}|z|^2)$ (see [9], sec. 1.1, Remark 1; for $n > 2$ one also has to use differentiation).

One can immediately check that

$$W(f_{m,n}) = f_{m+2,n} + nf_{m+1,n+1}.$$

Hence, from Theorem 1.2.1 we get the following formula for $g_{a,b}$.

Corollary 1.2.2. *For a pair of integers $a, b \geq 0$ of different parity one has*

$$g_{a,b} = \sum_{k \geq 0} k! \left(\binom{a}{k} + \binom{b}{k} \right) f_{a+b-k,k+1}$$

2. MINIMAL A_∞-ALGEBRA OF AN ELLIPTIC CURVE

2.1. **General construction.** Let us first recall the general construction of the A_∞-structure on the cohomology of a dg-algebra (A, d) equipped with a projector $\Pi : A \to B$ onto a subspace of $\ker(d)$ and a homotopy operator Q such that $1 - \Pi = dQ + Qd$. Merkulov's formula for this A_∞-structure (see [7]) was rewritten in [5] as a sum over trees:

$$m_n(b_1, \ldots, b_n) = -\sum_T \epsilon(T) m_T(b_1, \ldots, b_n).$$

Here T runs over all oriented planar rooted 3-valent trees with n leaves (different from the root) marked by $b_1, \ldots, b_n$ left to right, and the root marked by Π (we draw the tree in such a way that leaves are above, and every vertex has two edges coming from above and one from below). The expression $m_T(b_1, \ldots, b_n)$ is obtained by going down from leaves to the root, applying multiplication in A at every vertex and applying the operator Q at every inner edge (see [5], sec. 6.4, for details). The sign $\epsilon(T)$ has form

$$\epsilon(T) = \prod_v (-1)^{|e_1(v)|+(|e_2(v)|-1)\deg(e_1(v))},$$

where v runs through vertices of T, $(e_1(v), e_2(v))$ is the pair of edges above v, for an edge e we denote by $|e|$ the total number of leaves above e and by $\deg(e)$ the sum of degrees of all leaves above e (recall that leaves are marked by b_i).

Lemma 2.1.1. *Assume in addition that* $\Pi Q = Q\Pi = Q^2 = 0$. *Let* $(b_1, \ldots, b_n)$ *be a collection of elements in* B, *where* $n \geq 3$, *such that* $b_i = 1$ *for some* i. *Then* $m_n(b_1, \ldots, b_n) = 0$.

Proof. It is convenient to use Merkulov's original formula

$$m_n(b_1, \ldots, b_n) = \Pi\lambda_n(b_1, \ldots, b_n),$$

where $\lambda_n : A^{\otimes n} \to A$ are defined for $n \geq 2$ by the following recursion: $\lambda_2(a_1, a_2) = a_1 a_2$,

$$\lambda_n(a_1, \ldots, a_n) = \pm Q(\lambda_{n-1}(a_1, \ldots, a_{n-1})) \cdot a_n \pm a_1 \cdot Q(\lambda_{n-1}(a_2, \ldots, a_n)) + \sum_{k+l=n;k,l\geq 2} \pm Q(\lambda_k(a_1, \ldots, a_k)) \cdot Q(\lambda_l(a_{k+1}, \ldots, a_n)).$$

Since, $\Pi Q = 0$, it is enough to prove that $\lambda_n(b_1, \ldots, b_n) \in Q(A)$. Let us use induction in n. In the case $n = 3$ we have

$$\lambda_3(b_1, b_2, b_3) = Q(b_1 b_2) b_3 \pm b_1 Q(b_2 b_3)$$

and the assertion follows immediately from the fact that $Q(B) = 0$. Suppose now that $n \geq 4$ and the assertion holds for all $n' < n$. Since $Q^2 = 0$, the induction assumption easily implies that the first two terms in the recursive formula for λ_n belong to $Q(A)$. Similarly, all the remaining terms vanish if $n \geq 5$. In the case $n = 4$ the term $Q(b_1 b_2) \cdot Q(b_3 b_4)$ also vanishes because either $b_1 b_2 \in B$ or $b_3 b_4 \in B$ and $Q(B) = 0$. □

2.2. **The case of an elliptic curve.** Let $C = \mathbb{C}/(\mathbb{Z} \oplus \mathbb{Z}\tau)$ be a complex elliptic curve. We denote by L the holomorphic line bundle of degree 1 on C, such that the theta-function $\theta(z, \tau)$ descends to a global section of L. We consider the Dolbeault dg-algebra

$$A = (\Omega^{0,*} \otimes \mathcal{E}nd(\mathcal{O}_C \oplus L), \overline{\partial}).$$

Its cohomology B is the direct sum of the following components:
(i) $\mathrm{Hom}(\mathcal{O}, \mathcal{O})$ and $\mathrm{Hom}(L, L)$, both generated by identity maps;
(ii) $\mathrm{Hom}(\mathcal{O}, L)$, one-dimensional space;
(iii) $\mathrm{Ext}^1(L, \mathcal{O})$, one-dimensional space;
(iv) $\mathrm{Ext}^1(\mathcal{O}, \mathcal{O})$ and $\mathrm{Ext}^1(L, L)$, both isomorphic to the one-dimensional space $H^1(\mathcal{O})$.

To construct the homotopy operator Q, as in [8], we use the flat metric on C and the hermitian metric on L given by

$$(f,g)=\int_C f(z)\overline{g(z)}\exp(-2\pi\frac{\operatorname{Im}(z)^2}{\operatorname{Im}(\tau)})dxdy,$$

where $z=x+iy$. Then we set $Q=\overline{\partial}^*G$, where G is the Green operator corresponding to the Laplacian $\overline{\partial}^*\overline{\partial}+\overline{\partial}\overline{\partial}^*$. Then $B\subset A$ is exactly the space of harmonic forms, and $\Pi: A\to B$ is the orthogonal projection. Let us fix the following harmonic generators in the above components:
(i) $\mathrm{id}_{\mathcal{O}}$ and id_L;
(ii) $\theta=\theta(z,\tau)$ viewed as a holomorphic section of L;
(iii) $\eta:=\sqrt{2\operatorname{Im}(\tau)}\cdot\overline{\theta(z,\tau)}\exp(-2\pi\frac{\operatorname{Im}(z)}{\operatorname{Im}(\tau)^2})d\overline{z}$ viewed as a $(0,1)$-form with values in L^{-1};
(iv) $\xi=d\overline{z}$. When it is viewed as an element of $\operatorname{Ext}^1(L,L)$ we write ξ_L.

Note that we have a natural symmetric bilinear pairing on $A=A^0\oplus A^1$ given by

$$\langle\alpha,\beta\rangle=\frac{1}{2i\operatorname{Im}(\tau)}\cdot\int_C \operatorname{Tr}(\alpha\circ\beta)\wedge dz,$$

where α and β are homogeneous elements such that $\deg(\alpha)+\deg(\beta)=1$. The normalization is chosen in such a way that

$$\langle\xi,1\rangle=\frac{1}{2i\operatorname{Im}(\tau)}\cdot\int_C d\overline{z}\wedge dz=1.$$

By Serre duality, the induced pairing between B^0 and B^1 is nondegenerate. Also, by Theorem 1.1 of [8], the A_∞-structure on B satisfies the following cyclic symmetry:

$$\langle m_n(\alpha_1,\ldots,\alpha_n),\alpha_{n+1}\rangle=(-1)^{n(\deg(\alpha_1)+1)}\langle\alpha_1,m_n(\alpha_2,\ldots,\alpha_{n+1})\rangle. \tag{2.2.1}$$

The product m_2 on B is just the induced product on cohomology. The only interesting products are $m_2(\theta,\eta)\in\operatorname{Ext}^1(\mathcal{O},\mathcal{O})$ and $m_2(\eta,\theta)\in\operatorname{Ext}^1(L,L)$. Both are proportional to the generator ξ. To find the coefficient of proportionality it is enough to compute

$$\langle m_2(\theta,\eta),\mathrm{id}_{\mathcal{O}}\rangle=\langle m_2(\eta,\theta),\mathrm{id}_L\rangle=\frac{1}{2i\operatorname{Im}(\tau)}\cdot\int_C\theta\cdot\eta\wedge dz.$$

The above integral is well known:

$$\frac{\sqrt{2\operatorname{Im}(\tau)}}{2i\operatorname{Im}(\tau)}\cdot\int_C\theta(z,\tau)\overline{\theta(z,\tau)}\exp(-2\pi\frac{\operatorname{Im}(z)}{\operatorname{Im}(\tau)^2})d\overline{z}\wedge dz=1.$$

Thus, we obtain

$$m_2(\theta,\eta)=\xi,\quad m_2(\eta,\theta)=\xi_L. \tag{2.2.2}$$

By Lemma 2.1.1, every higher product m_n containing $\mathrm{id}_{\mathcal{O}}$ or id_L vanishes. Together with the cyclic symmetry (2.2.1) this implies that the only potentially nonzero higher products are of the following types:
(I) $m_n((\xi)^a,\theta,(\xi_L)^b,\eta,(\xi)^c,\theta,(\xi_L)^d)\in\operatorname{Hom}(\mathcal{O},L)$,
(II) $m_n((\xi_L)^a,\eta,(\xi)^b,\theta,(\xi_L)^c,\eta,(\xi)^d)\in\operatorname{Ext}^1(L,\mathcal{O})$,
(III) $m_n((\xi)^a,\theta,(\xi_L)^b,\eta,(\xi)^c,\theta,(\xi_L)^d,\eta,(\xi)^e)\in\operatorname{Hom}(\mathcal{O},\mathcal{O})$,
(IV) $m_n((\xi_L)^a,\eta,(\xi)^b,\theta,(\xi_L)^c,\eta,(\xi)^d,\theta,(\xi_L)^e)\in\operatorname{Hom}(L,L)$,

where we denote by $(\xi)^a$ the string $(\xi,\dots,\xi)$ with ξ repeated a times.

By the cyclic symmetry (2.2.1), we have

$$\langle m_n((\xi_L)^a,\eta,(\xi)^b,\theta,(\xi_L)^c,\eta,(\xi)^d),\theta\rangle = \langle m_n((\xi)^b,\theta,(\xi_L)^c,\eta,(\xi)^d,\theta,(\xi_L)^a),\eta\rangle,$$
$$\langle m_n((\xi)^a,\theta,(\xi_L)^b,\eta,(\xi)^c,\theta,(\xi_L)^d,\eta,(\xi)^e),\xi\rangle = \langle m_n((\xi)^{a+e+1},\theta,(\xi_L)^b,\eta,(\xi)^c,\theta,(\xi_L)^d),\eta\rangle,$$
$$\langle m_n((\xi_L)^a,\eta,(\xi)^b,\theta,(\xi_L)^c,\eta,(\xi)^d,\theta,(\xi_L)^e),\xi_L\rangle = \langle m_n((\xi)^b,\theta,(\xi_L)^c,\eta,(\xi)^d,\theta,(\xi_L)^{a+e+1}),\eta\rangle.$$

Hence, it is enough to compute the products of type (I), i.e., the coefficients

$$\langle m_n((\xi)^a,\theta,(\xi_L)^b,\eta,(\xi)^c,\theta,(\xi_L)^d),\eta\rangle.$$

2.3. **Calculation I: combinatorial part.** We start by computing some signs $\epsilon(T)$. For a pair of oriented planar rooted 3-valent trees T_1,T_2 let us denote by $\mathrm{join}(T_1,T_2)$ the tree (of the same type) obtained by joining together the roots of T_1 and T_2 and adding a root to the obtained new vertex (we keep T_1 on the left from T_2 in the plane).

Lemma 2.3.1. *Let $T=\mathrm{join}(T_1,T_2)$, where each T_i has n_i+1 leaves ($i=1,2$), exactly one of which is marked by a degree 0 element, and the rest marked by degree 1 elements. Assume also that in each T_i no two leaves of degree 1 can be attached to the same vertex. Then*

$$\epsilon(T)=(-1)^{\binom{n_1+n_2+2}{2}+n_2}.$$

Proof. By definition,

$$\epsilon(T)=\epsilon(T_1)\epsilon(T_2)\cdot(-1)^{(n_1+1)+n_1n_2},$$

so it remains to compute $\epsilon(T_i)$ for $i=1,2$. Note that under our assumptions each tree T_i has a very simple structure: it has the main stem from the root to the leaf of degree 0, to which leaves of degree 1 can attach on the left and on the right:



Suppose we have a pair of consecutive vertices v and w on the stem (with v above w) such that there is a degree 1 leaf attaching to v on the left and a degree 1 leaf attaching to w on the right. Let a be the number of degree 1 leaves above v. Then the contribution of v into the product defining $\epsilon(T_i)$ equals $(-1)^{a+1}$, while the contribution of w equals $(-1)^{a+2}$. Hence, the contribution of both v and w is -1. It is easy to check that if T' is the tree obtained from T by reversing the order of attaching these two leaves at vertices v and w then the contribution of these vertices into $\epsilon(T')$ will still be -1. Assume that T_i has exactly a (resp., b) leaves to the left (resp., to the right) of the degree 0 leaf, and let $v_1, \dots, v_a$ (resp., $w_1, \dots, w_b$) be the vertices on the stem to which they attach. By the above observation, it is enough to consider the case when all the vertices $v_1, \dots, v_a$ are above $w_1, \dots, w_b$:



Then one can easily calculate that

$$\epsilon(T_i) = (-1)^{\binom{a+b+1}{2}}.$$

Hence,

$$\epsilon(T) = (-1)^{\binom{n_1+1}{2}+\binom{n_2+1}{2}+(n_1+1)+n_1n_2} = (-1)^{\binom{n_1+n_2+2}{2}+n_2}.$$

□

Next, we consider the terms $m_T((\xi)^a, \theta, (\xi_L)^b, \eta, (\xi)^c, \theta, (\xi_L)^d)$. Note that if two leaves of degree 1 in T are attached to the same vertex then m_T vanishes because it will involve taking products of two elements of degree 1 in A. Henceforward, we assume that no two leaves of degree 1 can be attached to the same vertex. It is convenient to introduce the operator

$$H_L : C^\infty(L) \to C^\infty(L) : s \mapsto Q(s \cdot d\overline{z})$$

and a similar operator $H_{\mathcal{O}}$ on C^∞-functions. The next result follows immediately from the definition.

Lemma 2.3.2. *(i) Let $T = \mathrm{join}(T_1, T_2)$, where the leaves of T_1 are marked (from left to right) by*

$$(\xi)^a, \theta, (\xi_L)^b, \eta, (\xi)^{c_1},$$

while the leaves of T_2 are marked by

$$(\xi)^{c_2}, \theta, (\xi_L)^d.$$

Let $a = a_1 + a_2$, where a_1 leftmost leaves of T_1 get attached to the main stem of T_1 below the vertex where η attaches to the main stem, and the next a_2 leaves get attached to the stem above this vertex:



Then

$$m_T((\xi)^a, \theta, (\xi_L)^b, \eta, (\xi)^{c_1+c_2}, \theta, (\xi_L)^d) = \Pi\left([H_{\mathcal{O}}^{a_1+c_1} Q(H_L^{a_2+b}(\theta)\cdot\eta)]\cdot H_L^{c_2+d}(\theta)\right).$$

(ii) Let $T = \mathrm{join}(T_1, T_2)$, where the leaves of T_1 are marked by

$$(\xi)^a, \theta, (\xi_L)^{b_1},$$

while the leaves of T_2 are marked by

$$(\xi_L)^{b_2}, \eta, (\xi)^c, \theta, (\xi_L)^d.$$

Let $d = d_1 + d_2$, where exactly d_2 rightmost leaves of T_2 get attached to the main stem of T_2 below the vertex where η attaches to the main stem. Then

$$m_T((\xi)^a, \theta, (\xi_L)^{b_1+b_2}, \eta, (\xi)^c, \theta, (\xi_L)^d) = \Pi\left([H_{\mathcal{O}}^{b_2+d_2} Q(H_L^{c+d_1}(\theta)\cdot\eta)]\cdot H_L^{a+b_1}(\theta)\right).$$

Combining Lemmas 2.3.1 and 2.3.2 we arrive at the following expression for m_n.

Lemma 2.3.3. *One has*

$$\langle m_n((\xi)^a, \theta, (\xi_L)^b, \eta, (\xi)^c, \theta, (\xi_L)^d), \eta\rangle =$$
$$(-1)^{\binom{n}{2}+1}\cdot \sum_{a=a_1+a_2;\, c=c_1+c_2} \binom{a_2+b}{a_2}\binom{a_1+c_1}{a_1}\binom{c_2+d}{c_2}\phi(a_2+b, a_1+c_1, c_2+d) +$$
$$(-1)^{\binom{n}{2}+n+1}\cdot \sum_{b=b_1+b_2;\, d=d_1+d_2} \binom{c+d_1}{c}\binom{b_2+d_2}{b_2}\binom{a+b_1}{a}\phi(c+d_1, b_2+d_2, a+b_1),$$

where

$$\phi(m,n,p) = (-1)^p \langle \Pi\left([H^n_{\mathcal{O}} Q(H^m_L(\theta)\cdot\eta)]\cdot H^p_L(\theta)\right), \eta\rangle.$$

2.4. **Calculation II: analytic part.** We will use real coordinates u, v on $\mathbb{C}$ such that $z = u + v\tau$. Let us also set $a = \mathrm{Im}(\tau)$.

Lemma 2.4.1. *Consider the differential operator* $D = -\frac{a}{\pi}\frac{\partial}{\partial z} - 2iav$. *Then for* $k \geq 0$ *one has:*

$$H^k_L\theta = \frac{1}{k!}\cdot D^k\theta = \frac{(-2ia)^k}{k!}\cdot\sum_{n\in\mathbb{Z}}(n+v)^k\exp(\pi i\tau n^2 + 2\pi i n z), \tag{2.4.1}$$

$$\sqrt{2a}(D^k\theta)\cdot\overline{\theta}\exp(-2\pi a v^2) = \sum_{(m,n)\in\mathbb{Z}^2}(-1)^{mn}(m\overline{\tau}-n)^k\exp(-\frac{\pi}{2a}|m\tau-n|^2 + 2\pi i(mu+nv)). \tag{2.4.2}$$

Proof. The case $k = 0$ of the identity (2.4.2) is well-known (see [8], eq. (2.2)). The general case follows easily by applying D^k. Next, let us prove (2.4.1) by induction in k. Recall that the operator $Q : \Omega^{0,1}(L) \to \Omega^{0,0}(L)$ is uniquely determined by the following two properties: $\overline{\partial}\circ Q = \mathrm{id}$ and the image of Q is orthogonal to θ. Thus, $H^k_L\theta$ equals the unique function f such that $\frac{\partial f}{\partial\overline{z}} = \frac{D^{k-1}\theta}{(k-1)!}$ and $(f,\theta) = 0$. Thus, it is enough to check the identities

$$D^k\theta = (-2ia)^k\cdot\sum_{n\in\mathbb{Z}}(n+v)^k\exp(\pi i\tau n^2 + 2\pi i n z), \tag{2.4.3}$$

$$\frac{\partial}{\partial\overline{z}}D^k\theta = kD^{k-1}\theta, \tag{2.4.4}$$

$$(D^k\theta,\theta) = 0$$

for $k > 0$. The latter follows immediately from the Fourier expansion (2.4.2), since $(D^k\theta,\theta)$ is proportional to the Fourier coefficient of this expansion corresponding to $(m,n) = (0,0)$. To check (2.4.3) one can apply D to the similar expansion for $D^{k-1}\theta$. Finally, one checks (2.4.4) by applying $\frac{\partial}{\partial\overline{z}}$ to the right-hand side of (2.4.3). □

Now we can calculate the expressions $\phi(m,n,p)$ (see Lemma 2.3.3).

Lemma 2.4.2. *One has*

$$\phi(k,l,p) = \frac{1}{k!p!}(\frac{a}{\pi})^{l+1}\cdot\sum_{\omega\in\mathbb{Z}+\mathbb{Z}\tau\setminus\{0\}}\frac{\overline{\omega}^{k+p}}{\omega^{l+1}}\exp(-\frac{\pi}{a}|\omega|^2) = \frac{1}{k!p!}(\frac{a}{\pi})^{k+l+p+1}f_{k+p,l+1}(\mathbb{Z}+\mathbb{Z}\tau).$$

Proof. It is easy to see that the operator $Q : \Omega^{0,1}\to\Omega^{0,0}$ is given by

$$Q(\exp(2\pi i(mu+nv))d\overline{z}) = \begin{cases}\frac{a}{\pi(m\tau-n)}\exp(2\pi i(mu+nv)), & (m,n)\neq(0,0),\\ 0 & (m,n) = (0,0).\end{cases}$$

Hence, using Lemma 2.4.1 we obtain

$$Q(H_L^k(\theta)\cdot\eta)=\frac{1}{k!}Q(\sqrt{2a}(D^k\theta)\cdot\overline{\theta}\exp(-2\pi a v^2)=$$
$$\frac{a}{k!\pi}\cdot\sum_{(m,n)\in\mathbb{Z}^2\setminus\{(0,0)\}}(-1)^{mn}\frac{(m\overline{\tau}-n)^k}{(m\tau-n)}\exp(-\frac{\pi}{2a}|m\tau-n|^2+2\pi i(mu+nv)).$$

Therefore,
(2.4.5)
$$H_{\mathcal{O}}^lQ(H_L^k(\theta)\cdot\eta)=\frac{1}{k!}(\frac{a}{\pi})^{l+1}\cdot\sum_{(m,n)\in\mathbb{Z}^2\setminus\{(0,0)\}}(-1)^{mn}\frac{(m\overline{\tau}-n)^k}{(m\tau-n)^{l+1}}\exp(-\frac{\pi}{2a}|m\tau-n|^2+2\pi i(mu+nv)).$$

Next, comparing the formulas for η and for the metric on L we observe that for a C^∞-section f of L one has $(f,\theta)=0$ if and only if $\langle f,\eta\rangle=0$. Hence, for $f\in\mathbb{C}^\infty(L)$ one has

$$\langle\Pi(f),\eta\rangle=\langle f,\eta\rangle$$

(since Π is the orthogonal projection onto $\mathbb{C}\theta$). Therefore,

$$\phi(k,l,p)=(-1)^p\langle[H_{\mathcal{O}}^lQ(H_L^k(\theta)\cdot\eta]\cdot H_L^p(\theta),\eta\rangle=(-1)^p\langle H_{\mathcal{O}}^lQ(H_L^k(\theta)\cdot\eta,H_L^p(\theta)\cdot\eta\rangle.$$

Now the right-hand side can be computed using the Fourier expansion for $H_L^p(\theta)$ from Lemma 2.4.1 and the Fourier expansion (2.4.5):

$$\phi(k,l,p)=\frac{1}{k!p!}(\frac{a}{\pi})^{l+1}\cdot\sum_{(m,n)\in\mathbb{Z}^2\setminus\{(0,0)\}}\frac{(m\overline{\tau}-n)^{k+p}}{(m\tau-n)^{l+1}}\exp(-\frac{\pi}{a}|m\tau-n|^2).$$

□

2.5. **Calculation III: conclusion.** It remains to put everything together. Substituting the expressions for $\phi(k,l,p)$ found in Lemma 2.4.2 into the formula of Lemma 2.3.3 we get

$$\langle m_n((\xi)^a,\theta,(\xi_L)^b,\eta,(\xi)^c,\theta,(\xi_L)^d),\eta\rangle=$$
$$(-1)^{\binom{n}{2}+1}(\frac{\operatorname{Im}(\tau)}{\pi})^{n-2}\cdot\sum_{a=a_1+a_2;c=c_1+c_2}\frac{(a_1+c_1)!}{a_1!a_2!c_1!c_2!b!d!}f_{a_2+c_2+b+d,a_1+c_1+1}+$$
$$(-1)^{\binom{n}{2}+n+1}(\frac{\operatorname{Im}(\tau)}{\pi})^{n-2}\cdot\sum_{b=b_1+b_2;d=d_1+d_2}\frac{(b_2+d_2)!}{b_1!b_2!d_1!d_2!a!c!}f_{b_1+d_1+a+c,b_2+d_2+1}.$$

Since $f_{k,l}=0$ unless $k+l$ is even, this immediately implies that $m_n=0$ for odd n. Now assuming that n is even we can rewrite the above equation as follows (denoting $k=a_1+c_1$ and $l=b_2+d_2$):

$$(-1)^{\binom{n}{2}+1}(\frac{\pi}{\operatorname{Im}(\tau)})^{n-2}\cdot\langle m_n((\xi)^a,\theta,(\xi_L)^b,\eta,(\xi)^c,\theta,(\xi_L)^d),\eta\rangle=$$
$$\frac{1}{b!d!}\cdot\sum_{k\geq 0}C(a,c,k)f_{a+c-k+b+d,k+1}+\frac{1}{a!c!}\cdot\sum_{l\geq 0}C(b,d,l)f_{b+d-l+a+c,l+1},$$

where

$$C(a,c,k)=\sum_{a_1+c_1=k;a_1\le a,c_1\le c}\frac{k!}{a_1!c_1!(a-a_1)!(c-c_1)!}=\frac{k!}{a!c!}\sum_{a_1+c_1=k}\binom{a}{a_1}\binom{c}{c_1}=\frac{k!}{a!c!}\binom{a+c}{k}$$

Thus, our formula for $\langle m_n((\xi)^a,\theta,(\xi_L)^b,\eta,(\xi)^c,\theta,(\xi_L)^d),\eta\rangle$ takes form

$$(-1)^{\binom{n}{2}+1}(\frac{\pi}{\operatorname{Im}(\tau)})^{n-2}\cdot\langle m_n((\xi)^a,\theta,(\xi_L)^b,\eta,(\xi)^c,\theta,(\xi_L)^d),\eta\rangle=$$

$$\frac{1}{a!b!c!d!}\cdot\sum_{k\ge 0}k!\left(\binom{a+c}{k}+\binom{b+d}{k}\right)f_{n-3-k,k+1}.$$

Taking into account Corollary 1.2.2 we obtain for even n

$$\langle m_n((\xi)^a,\theta,(\xi_L)^b,\eta,(\xi)^c,\theta,(\xi_L)^d),\eta\rangle=(-1)^{\binom{n}{2}+1}\frac{1}{a!b!c!d!}\cdot(\frac{\operatorname{Im}(\tau)}{\pi})^{n-2}\cdot g_{a+c,b+d}. \tag{2.5.1}$$

Let us summarize our calculations. We set

$$M(a,b,c,d):=(-1)^{\binom{a+b+c+d+1}{2}}\frac{1}{a!b!c!d!}\cdot(\frac{\operatorname{Im}(\tau)}{\pi})^{a+b+c+d+1}\cdot g_{a+c,b+d}. \tag{2.5.2}$$

Theorem 2.5.1. *The only non-trivial higher products m_n of the A_∞-structure on $B=\operatorname{Ext}^*(\mathcal{O}\oplus L,\mathcal{O}\oplus L)$ are of the form*

$$m_n((\xi)^a,\theta,(\xi_L)^b,\eta,(\xi)^c,\theta,(\xi_L)^d)=M(a,b,c,d)\cdot\theta,$$

$$m_n((\xi_L)^a,\eta,(\xi)^b,\theta,(\xi_L)^c,\eta,(\xi)^d)=M(a,b,c,d)\cdot\eta,$$

$$m_n((\xi)^a,\theta,(\xi_L)^b,\eta,(\xi)^c,\theta,(\xi_L)^d,\eta,(\xi)^e)=M(a+e+1,b,c,d)\cdot\mathrm{id}_{\mathcal{O}},$$

$$m_n((\xi_L)^a,\eta,(\xi)^b,\theta,(\xi_L)^c,\eta,(\xi)^d,\theta,(\xi_L)^e)=M(a+e+1,b,c,d)\cdot\mathrm{id}_L\,.$$

All products m_n with odd n vanish.

Remarks. 1. Assume that τ belongs to a ring of integers of an imaginary quadratic field (so that our elliptic curve admits complex multiplication). Set $\varpi=2\pi|\eta(q)|^2$, where $\eta(q)$ is the Dedekind's η-function, and $q=\exp(2\pi i\tau)$. Then the numbers $\varpi^{-a-b-1}\cdot g_{a,b}$ are algebraic over $\mathbb{Q}$ (see [10], sec. VI.6). Hence, if we multiply our basis elements of degree 1 (η, ξ and ξ_L) by the factor $|\eta(q)|^{-2}$ then the structure constants of our A_∞-structure with respect to the new basis will be algebraic over $\mathbb{Q}$.
2. Another meaningful rescaling is obtained if we multiply our basis elements of degree 1 by the factor $\pi/\operatorname{Im}(\tau)$. Then the new structure constants $M'(a,b,c,d)$ will all have limit at the cusp $\operatorname{Im}(\tau)\to+\infty$. Namely, using the well known q-series for the Eisenstein series, one can easily check that the only nonzero limiting values will be

$$M'(i,0,j,0)=M'(0,i,0,j)\to(-1)^{\binom{i+j+1}{2}}\cdot\binom{i+j}{i}\cdot 2\zeta(i+j+1),$$

where $i+j$ is odd.

2.6. **A_∞-constraint.** The A_∞-axiom (we follow [4], 3.1 for sign conventions) gives certain quadratic equations on coefficients $(M(a,b,c,d))$ and hence leads to identities for $g_{m,n}$. For example, applying this axiom to the string

$$(\xi)^a, \theta, (\xi_L)^b, \eta, (\xi)^c, \theta, (\xi_L)^d, \eta, (\xi)^e, \theta, (\xi_L)^f,$$

where a,b,c,d,e,f are positive, we get

$$\begin{aligned}
&\sum_{a=a_1+a_2;d=d_1+d_2} (-1)^{(a_2+b+c+d_1+1)(a_1+d_2+e+f)+a_1} M(a_2,b,c,d_1)M(a_1,d_2,e,f)+\\
&\sum_{b=b_1+b_2;e=e_1+e_2} (-1)^{(b_2+c+d+e_1+1)(a+b_1+e_2+f+1)+a+b_1+1} M(b_2,c,d,e_1)M(a,b_1,e_2,f)+\\
&\sum_{c=c_1+c_2;f=f_1+f_2} (-1)^{(c_2+d+e+f_1+1)(a+b+c_1+1+f_2)+a+b+c_1} M(c_2,d,e,f_1)M(a,b,c_1,f_2)=0.
\end{aligned}$$

When one of a,b,c,d,e,f is zero, additional terms will arise due to the presence of double products. For example, for the string

$$(\xi)^a, \theta, \eta, \theta, \eta, \theta, (\xi_L)^b$$

we get

$$\begin{aligned}
&\sum_{a=a_1+a_2} (-1)^{(a_2+1)(a_1+b)+a_1} M(a_2,0,0,0)M(a_1,0,0,b)+\\
&\sum_{b=b_1+b_2} (-1)^{(b_1+1)(a+b_2+1)+a} M(0,0,0,b_1)M(a,0,0,b_2)+\\
&(-1)^a[M(a+1,0,0,b)-M(a,1,0,b)+M(a,0,1,b)-M(a,0,0,b+1)]+\\
&\delta_{b,0}(-1)^a M(a+1,0,0,0)-\delta_{a,0}M(b+1,0,0,0)=0.
\end{aligned}$$

Substituting into the above identities the expressions for $M(a,b,c,d)$ from (2.5.2) we arrive at the following result.

Proposition 2.6.1. *(i) For positive integers a,b,c,d,e,f one has*

$$\begin{aligned}
&\sum_{a=a_1+a_2;d=d_1+d_2} (-1)^{c+d_1+1}\binom{a}{a_1}\binom{d}{d_1} g_{a_2+c,b+d_1}g_{a_1+e,d_2+f}+\\
&\sum_{b=b_1+b_2;e=e_1+e_2} (-1)^{b_2+1}\binom{b}{b_1}\binom{e}{e_1} g_{b_2+d,c+e_1}g_{a+e_2,b_1+f}+\\
&\sum_{c=c_1+c_2;f=f_1+f_2} (-1)^{c_1}\binom{c}{c_1}\binom{f}{f_1} g_{c_2+e,d+f_1}g_{a+c_1,b+f_2}=0.
\end{aligned}$$

(ii) For integers $a,b\geq 0$ one has

$$\sum_{a=a_1+a_2}\binom{a}{a_1}g_{a_1,0}g_{a_2,b}-\frac{a+2+\delta_{b,0}}{a+1}g_{a+1,b}=\sum_{b=b_1+b_2}\binom{b}{b_1}g_{0,b_1}g_{a,b_2}-\frac{b+2+\delta_{a,0}}{b+1}g_{a,b+1}.$$

Note that the identity in (ii) gives a recursive formula for $g_{a+1,b}$ in terms of all $g_{a',b'}$ with $a' \leq a$. Since $g_{0,n} = n!e^*_{n+1}$, we recover the fact that all $g_{a,b}$ are polynomials in (e^*_n) with rational coefficients. For example, in the case $a = 0$ the obtained identity (for even n)

$$2g_{1,n} = -\sum_{n=m+k} \binom{n}{m} g_{0,m}g_{0,k} + \frac{n+3}{n+1} g_{n+1,0}$$

is equivalent to the formula

$$\frac{1}{n} We^*_n = -\sum_{n=m+k} e^*_{m+1}e^*_{k+1} + (n+3)e_{n+2}$$

found in [10], VI.5.

Bibliography

[1] A. Bondal, D. Orlov, *Derived categories of coherent sheaves*, Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002), 47–56, Higher Ed. Press, Beijing, 2002.

[2] N. Katz, *p-adic interpolation of real analytic Eisenstein series*, Annals of Math. 104 (1976), 459–571.

[3] B. Keller, *Introduction to A-infinity algebras and modules*, Homology Homotopy Appl. 3 (2001), no. 1, 1–35.

[4] B. Keller, *A-infinity algebras, modules and functor categories*, in *Trends in representation theory of algebras and related topics*, 67–93, AMS, Providence, RI, 2006.

[5] M. Kontsevich, Y. Soibelman, *Homological mirror symmetry and torus fibration*, in *Symplectic geometry and mirror symmetry (Seoul, 2000)*, 203–263, World Sci. Publishing, River Edge, NJ, 2001.

[6] K. Lefèvre-Hasegawa, *Sur les A_∞-catégories*, Thèse de doctorat, Université Denis Diderot – Paris 7, 2003, available at B. Keller's homepage.

[7] S. Merkulov, *Strong homotopy algebras of a Kähler manifold*, Internat. Math. Res. Notices 1999, no.3, 153–164.

[8] A. Polishchuk, *Homological mirror symmetry with higher products*, in *Proceedings of the Winter School on Mirror Symmetry, Vector Bundles and Lagrangian Submanifolds*, 247–259, AMS and International Press, 2001.

[9] A. Polishchuk, *Rapidly converging series for the Weierstrass zeta-function and for the Kronecker function*, Math. Research Letters 7 (2000), 493–502.

[10] A. Weil, *Elliptic functions according to Eisenstein and Kronecker.* Springer-Verlag, 1976.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OR 97403
E-mail address: `apolish@uoregon.edu`